


Cornell University
College of Veterinary Medicine



PURINA.




Boehringer
Ingelheim



Clinical Investigators' Day

2024





Cornell University College of Veterinary Medicine

Welcome to the 2024 Clinical Investigators' Day, sponsored by the Cornell University College of Veterinary Medicine. The primary goal of this forum is to provide an opportunity for residents and interns to showcase ongoing investigations carried out at Cornell University College of Veterinary Medicine. It is our hope that greater insights will be gained in the breadth and depth of clinical investigations conducted at the College and will serve as a catalyst to promote greater interactions among colleagues with clinical and basic science research interests.

Organizing Committee

Dr. Robert Goggs, Co-Chair
Dr. Kelly Hume, Co-Chair
Dr. Elizabeth Moore, Co-Chair
Dr. Tracy Stokol, Co-Chair
Mr. Doug Fink

The organizing committee thanks the following individuals who contributed to the success of the Day:

Dr. Manuel Martin-Flores
Mr. Dave Frank
Mr. Drew Kirby
Dr. John Parker
Dean Lorin Warnick

Images: iStock Photos and Drew Kirby



Disclaimer: The information presented in this publication comes exclusively from the contributing authors and their mentors identified in each abstract. These abstracts are preliminary and should not be cited in bibliographies. Material contained herein may be treated as personal communication and should be cited as such only with the consent of the author.

Sponsors

We thank the following sponsors for their generous support and commitment to Clinical Investigators' Day:

Corporate Sponsors



Nestlé PURINA



College of Veterinary Medicine

Animal Health Diagnostic Center (AHDC)

Baker Institute for Animal Health

Cornell University Hospital for Animals (CUHA)

Department of Biomedical Sciences

Department of Clinical Sciences

Department of Microbiology and Immunology

Department of Molecular Medicine

Department of Population Medicine and Diagnostic Sciences

Feline Health Center

College of Veterinary Medicine (cont.)

Office of the Dean

Office of Research and Graduate Education

Richard P. Riney Canine Health Center

University Programs

Center for Animal Resources and Education (CARE) at Cornell University



Schedule (cont.)

- **AMBULATORY SACROCOCCYGEAL EPIDURAL INJECTION PROVIDES SUPERIOR POST-OPERATIVE ANALGESIA AFTER FELINE OVARIOHYSTERECTOMY (OVH)**

Shana Wong – Anesthesiology and Pain Management Resident

Pg. 8

12:00 pm – 1:30 pm

Lunch

1:30 pm – 2:30 pm

Resident Presentations and Proposals – Moderated by Kimaya Bakhle (combined DVM/PhD student)

- **COMPARISON OF TWO LOCATIONS FOR EXTERNAL JUGULAR VEIN CANNULATION IN A PIG MODEL – PROPOSAL**

Rachel Mandelbaum – Anesthesiology and Pain Management Resident

Pg. 9

- **CHARACTERIZATION AND ASSESSMENT OF BIOLOGIC BEHAVIOR OF CANINE SPLENIC STROMAL SARCOMAS - PROPOSAL**

Corene Bruhns – Medical Oncology Resident

Pg. 10

- **ENHANCING POSTOPERATIVE RECOVERY IN RATS (RATTUS NORVEGICUS) AGAINST EXTENDED-RELEASE BUPRENORPHINE SIDE EFFECTS – PROPOSAL**

Dahihm Kim – Laboratory Animal Medicine Resident

Pg. 11

- **REFINEMENT OF CONTINUOUS POSITIVE AIRWAY PRESSURE (CPAP) THERAPY IN DOGS – PROPOSAL**

Victoria Albano – Anesthesiology and Pain Management Resident

Pg. 12


2:45 pm – 3:00 pm

Award Presentations

Dr. Lorin Warnick, Dean, College of Veterinary Medicine



Keynote Speaker



Mariana Diel de Amorim, DVM, DVSc, PhD, DACT

Assistant Professor of Theriogenology, Cornell University Department of Clinical Sciences

Dr. Diel de Amorim earned her DVM from Federal Rural University of Pernambuco, Brazil and completed both her PhD and residency in Theriogenology from University of Guelph, Canada. Her research interest is on equine theriogenology, especially pertaining to equine maternal recognition of pregnancy and equine endometritis. She has focused her efforts on looking at the molecular level on the endometrium, corpus luteum and embryo interaction.

2024 Judges



Doug Antczak, , VMD, PhD

Dorothy Havemeyer McConville Professor of Equine Medicine, Department of Microbiology and Immunology, Cornell University

Dr. Antczak's research program is focused on the health of horses. Through the Baker Institute's Equine Genetics Center, the Antczak laboratory has a long history of advancing basic knowledge and applying that knowledge in equine genetics, immunology, and reproduction. The cord that links these three distinct areas of science together is the intimate relationship between mother and fetus during pregnancy in mammals.



Galina Hayes, BVSc, DVSc, PhD, DACVECC, DACVS

Associate Professor of Small Animal Surgery, Department of Clinical Sciences, Cornell University

Dr. Hayes' research interests include the Surgical Techniques Initiative: Clinical trials evaluating changes in surgical methods that can reduce incisional pain and inflammation, reduce infection rates and improve healing in canine surgery. Other areas of interest include predictive scoring systems in clinical medicine, minimally invasive surgery, and trauma management.

2023 Judges (cont.)



Margaret McEntee, DVM, DACVIM, DACVR

Professor of Medical and Radiation Oncology, Department of Clinical Sciences, Cornell University

Dr. McEntee earned her DVM from Cornell University and completed residencies at North Carolina State University in both Medical Oncology and Radiation Oncology. Her research interests include Contrast Enhanced CT scanning, advancing the technical aspects of radiation therapy, and linear accelerator/3D treatment planning.



Peter V. Scrivani, DVM, DACVR

Professor and Section Chief of Diagnostic Imaging, Department of Clinical Sciences, Cornell University

Dr. Scrivani earned his DVM from Cornell University and completed his residency at The Ohio State University. His research uses imaging and epidemiologic techniques for investigating and establishing the best clinical practices in animal health care. His specific research focus is advancing the understanding of head and neck disorders, with emphasis on the role of the underlying morphology.



Moderators



Kimaya Bakhle

Combined DVM/PhD Student, College of Veterinary Medicine, Cornell University

Kimaya worked at a small animal clinic during high school, which sparked her interest in veterinary medicine. She pursued this interest through a pre-veterinary biochemistry program at Purdue University and is now enrolled in the Cornell Combined DVM/PhD Program. In the future, Kimaya hopes to use her background in veterinary medicine and cancer biology research to improve outcomes for animals and humans with cancer.



Megan Fahey

Combined DVM/PhD Student, College of Veterinary Medicine, Cornell University

Meg is a fifth-year Combined DVM/PhD student currently working in Dr. Gerlinde Van de Walle's lab. Her research focuses on the secreted products (i.e., secretome) of mesenchymal stromal cells and their potential as an anti-microbial and pro-regenerative therapeutic. Meg is passionate about translational research and aspires to be a clinician-scientist who studies stem cell-derived therapeutics for humans and veterinary species alike.



Stephanie Hon, DVM, DACVAA

Assistant Professor, Section of Anesthesiology and Pain Medicine, College of Veterinary Medicine, Cornell University

Dr. Hon completed her veterinary training at the University of Florida, her rotating internship at Rood and Riddle Equine Hospital, and her anesthesia residency at Cornell University. Her research interests include locoregional and vascular access techniques in pigs, with an emphasis on supporting translational research efforts and bringing those skills and improvements to the clinic.



Pamela S. Velarde, DVM

pv245@cornell.edu

Institution and Location

Universidad Científica del Sur, Lima, Peru
Peterson and Smith Equine Hospital, Ocala, FL
Hagyard Equine Medical Institute, Lexington, KY
Cornell University, Ithaca, New York

Degree

DVM
Internship
Fellowship
Residency

Year

2008
2021
2022
2022-Present

Current Position

2nd Year Resident, Large Animal Internal Medicine

Abstract Title:

ANTIMICROBIAL PRESCRIBING IN AN EQUINE TEACHING HOSPITAL

Authors Names:

Pamela S. Velarde¹, Kevin J. Cummings², Charisse Lapointe¹, Mariajose Cervantes¹, Sadie Hubler¹, Gillian A. Perkins¹

¹Department of Clinical Sciences, Cornell University, Ithaca, New York

²Department of Public and Ecosystem Health, Cornell University, Ithaca, New York

Project Mentor:

Mentor: Gillian A. Perkins DVM DACVIM, Department of Clinical Sciences, gap7@cornell.edu

Co-Mentor: Kevin J. Cummings DVM PhD, Department of Public and Ecosystem Health, kjc39@cornell.edu

Abstract:

Antimicrobial resistance is a global veterinary and human health concern. There is potential for bacteria to become resistant whenever an antimicrobial is administered, and therefore veterinarians should practice judicious use. Recent studies have shown that equine practitioners may over-prescribe antimicrobials, and antimicrobials classified as critically important by the World Health Organization (WHO) may be given when a first-line antimicrobial would be sufficient. The objective of this retrospective study was to describe the antimicrobial prescribing practices in an equine teaching hospital over a one-year period to highlight possible areas of improvement in antimicrobial stewardship. Electronic medical records from the Cornell University Equine Hospital were searched from January 1st to December 31st, 2021, data collected, and descriptive statistics performed. A total of 33.5% (543/1622) of equine patients were prescribed systemic antimicrobials in 2021. The most common antimicrobial drug classes used were penicillins and aminoglycosides. Most horses (75%) were prescribed ≤ 2 antimicrobials. About 24% of non-surgical patients were given antimicrobials when there was no evidence of infection on clinical exam and diagnostic testing. Surgical patients accounted for around 58% of those receiving antibiotics and many (about 33%) were clean surgeries. Cultures were only submitted about 16% of the time. In conclusion, antimicrobial prescribing practices could be improved at the equine teaching hospital.



Emilia Chrostek, DVM

ec798@cornell.edu

Institution and Location

Colorado State University
VCA Canada Calgary Animal Referral and Emergency
Centre, Calgary, Alberta, Canada
Cornell University, Ithaca, New York

Degree

DVM

Year

2021

Internship

2019

Residency

2021-Present

Current Position

2nd Year Resident in Dentistry and Oral Surgery

Abstract Title:

MORPHOLOGICAL STUDY OF PULP CAVITY ANATOMY OF CANINE TEETH IN DOMESTIC CATS USING MICRO-COMPUTED TOMOGRAPHY

Authors Names:

Emilia Chrostek¹, Santiago Peralta¹, Nadine Fiani¹

¹Department of Clinical Sciences, Cornell University, Ithaca, NY, USA


Project Mentor(s):

Mentor: Nadine Fiani, BVSc, Dipl. AVDC, FF-AVDC-OMFS, AVDC-ZWD, Department of Clinical Sciences, nf97@cornell.edu

Co-mentor: Santiago Peralta, DVM, Dipl. AVDC, FF-AVDC-OMFS, Department of Clinical Sciences, sp888@cornell.edu

Abstract:

An understanding of the pulp cavity anatomy of individual teeth is essential for success during endodontic therapy. The primary objective of this study was to document pulp cavity anatomy and summarize numerical data of maxillary and mandibular canine teeth of domestic cats using micro-computed tomography (micro-CT). Thirty-nine canine teeth from eleven domestic cat cadaveric specimens were extracted and scanned. Segmentation of the pulp cavity was performed using the Avizo (v2022.2) software. The morphological features of the pulp cavity including configuration, presence of apical deltas and lateral canals was recorded. A quantitative analysis was performed on thirty-one teeth to explore associations between variables using a linear mixed model. Correlation between pertinent continuous variables was assessed using a Pearson’s correlation test. Most pulp cavities exhibited a cylindrical configuration in the coronal third to an ovoid configuration in the middle to apical third. A ribbon-like canal was observed in 6/31 teeth (19%). All teeth depicted an apical delta, except for two teeth that showed a single canal exiting at the apex. In 15/31 teeth (48%), the primary root canal within the apical delta could be clearly identified and in 16/31 (52%) the primary root canal was indiscernible. Additionally, the pulp cavities of maxillary canine teeth were significantly larger and longer and the cusp-to-tip length was longer, when compared to mandibular teeth. The apical delta length was negatively correlated to pulp cavity volume. This study revealed that the canine tooth pulp cavity in cats varies considerably and should be a consideration during endodontic therapy.



Hanna R.M. Sfraga, DVM

hs783@cornell.edu

Institution and Location

Colorado State University, Ft. Collins, Colorado
Cornell University, Ithaca, New York

Degree

DVM
Residency

Year

2021
2022-Present

Current Position

2nd Year Resident in Large Animal Internal medicine

Abstract Title:

POST-MORTEM CHARACTERIZATION OF SUBCLINICAL INFLAMMATION OF NUCHAL BURSAE AND NUCHAL LIGAMENTS IN HORSES

Authors Names:

Hanna RM Sfraga¹, Elena A Demeter², Toby L Pinn-Woodcock², Cassandra Guarino², Marta Cercone¹

¹Department of Clinical Sciences, Cornell University, Ithaca, New York

²Department of Population Medicine and Diagnostic Sciences, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Marta Cercone, DVM, PhD, DACVIM-LA, Department of Clinical Sciences, mc957@cornell.edu

Abstract:

Nuchal bursitis is an uncommon and challenging cause of pain and decreased performance in horses. Radiographic identification of nuchal bursae and/or ligament mineralization is often considered an incidental finding in the absence of swelling and pain. Historically, the etiologic agent of main concern in septic bursitis was *Brucella abortus/suis*. Recently, increasing cases of nuchal bursitis associated with *Borrelia burgdorferi* infection have been reported.

The aims of this prospective study were to investigate the incidence of subclinical nuchal bursitis, characterize its histopathologic features and its association with *Borrelia burgdorferi*.

This cadaver study used a convenient population of horses in a *B. burgdorferi* endemic region undergoing euthanasia and tissue donation without history or clinical signs of nuchal bursitis and/or ligament pathology.

Fifteen horses, 4-29 years old, (5 geldings, 10 mares, various breeds) were enrolled. Serum and post-mortem nuchal bursae, synovial fluid and nuchal ligament were collected. The bursae and ligament were evaluated via histopathology score system, grading inflammation, edema, fibrosis, and mineralization (0-normal to 18-severe). Lyme Multiplex assay on serum and *Borrelia burgdorferi* PCR on bursae were performed.

Inflammation of the bursa was detected in four horses, and six more showed other mild changes (edema, fibrosis, mineralization). The overall histopathology score ranged from 1 to 7 (mean 2 ± 1.65). Five horses had positive Lyme Multiplex assay antibody titers, while all tissue samples were negative on *B. burgdorferi* PCR.

Our study suggests a high incidence of mild/moderate nuchal bursae and ligament histopathologic changes (67%), warranting further investigation of their causes and implications.



Elizabeth S. Lavin, DVM

esl92@cornell.edu

Institution and Location

Cornell University, Ithaca, New York
Cornell University, Ithaca, New York

Degree

DVM
Residency

Year

2021
2021-Present

Current Position

3rd Year Resident in Laboratory Animal Medicine

Abstract Title:

VENIPUNCTURE SITE INFLUENCES BLOOD DROP VOLUME IN C57BL/6 MICE

Authors Names:

Elizabeth S Lavin¹, Erica R Feldman¹, Scott M Soprano¹, Elizabeth S Moore^{1,2}

¹Center for Animal Resources and Education, Cornell University, Ithaca, New York

²Meinig School of Biomedical Engineering, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Elizabeth Moore, DVM, PhD, Center for Animal Resources and Education, esm84@cornell.edu

Abstract:

Many experiments require the collection of blood samples from mice. However, the size of mice limits the volume that may be safely collected. In IACUC protocols, investigators may report the amount of blood they collect from mice as a number of drops. Many institutions use an anecdotal conversion factor (1 blood drop = 25 μ l) to ensure that research protocols remain in compliance with institutional guidelines. Previous work has not experimentally determined the volume of a murine blood drop. In this 10-week crossover experiment, two phlebotomists bled thirty C57BL/6J mice from three sites (facial, saphenous, and tail). Male and female mice were weighed weekly and divided amongst 5 groups (n=6): left and right tail vein, left and right saphenous vein, and facial vein. A single blood drop from each site was weighed, and the volume of each drop was calculated using the average blood density determined from 8 mice bled at the end of the study. Site and side significantly influenced drop weight and thus calculated volume. Facial venipuncture produced the largest drop volume (mean: 21.7 μ l), followed by the saphenous (mean: 9.97 μ l) and tail (mean: 4.96 μ l) sites. Left-sided venipuncture was associated with slightly larger-volume drops, though effect size was small. The results of this study may be useful in more accurately estimating blood loss via conversion of drops to volume. Additionally, collection from saphenous and tail veins apparently minimizes blood loss relative to facial venipuncture and may optimize serial collection of small-volume samples and animal welfare.



Colin Chik, DVM

cpc77@cornell.edu

Institution and Location

Cornell University, Ithaca, New York
Cornell University, Ithaca, New York

Degree

DVM
Residency

Year

2020
2021-Present

Current Position

3rd Year Resident in Small Animal Surgery

Abstract Title:

INVESTIGATION OF THE EFFECTS OF PRE-STRETCHING ON LAPAROSCOPIC WORKING SPACE IN DOGS

Authors Names:

Chik C¹, Hayes G¹, Martin-Flores M¹, Buote NJ¹


¹Department of Clinical Sciences, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Nicole Buote DVM DACVS-SA (MIS Founding Fellow), Department of Clinical Science

Abstract:

Laparoscopy is increasingly used in veterinary patients because it decreases surgical trauma, improves cosmetic appearance, and has fewer surgical complications such as hemorrhage, dehiscence, and infection. Standard laparoscopy uses carbon dioxide (CO₂) insufflation to expand the abdomen, allowing for visualization and introduction of instrumentation. An inability to achieve adequate working space during laparoscopy can result in unnecessary tissue trauma, surgical error, prolonged surgical times, and the need to convert to an open procedure. Several methods can improve the surgical working space. Pre-stretching of the abdominal wall has shown the most promise in human and porcine studies. The goal of our prospective study was to determine if transient pre-stretching of the abdominal wall would provide improved working space during canine elective laparoscopic procedures. In fifty healthy client-owned dogs, we pre-stretched the abdominal wall prior to laparoscopy by insufflating to an intraabdominal pressure of 10 mmHg for three minutes. We then carried out the laparoscopic procedure at a desufflated pressure of 6 mmHg. We measured changes in skin and intraabdominal working length every 30 minutes intra-operatively. We found a median increase in intra-abdominal working length of 4.44% (IQR 5.74%) and intra-abdominal working volume of 6.92% (IQR 5.36%) after abdominal pre-stretching compared to initial 6 mmHg IAP measurements ($p < 0.001$). Distance measurements at the end of surgery were similar to those obtained immediately after pre-stretching. In conclusion, pre-stretching of the abdominal wall improved working space and volume and the effects of pre-stretching persisted throughout the laparoscopic procedure.



Kailah M. Buchanan, DVM

kmb386@cornell.edu

Institution and Location

St. George’s University, Grenada
University of Illinois, Urbana, Illinois
Kansas State University, Manhattan, Kansas
Cornell University, Ithaca, New York

Degree

DVM
Internship
Internship
Internship

Year

2021
2022
2023
2023-Present

Current Position

Intern in Cardiology

Abstract Title:

CHANGES TO PACEMAKER PROGRAMMING PARAMETERS OVER TIME IN DOGS

Authors Names:

Kailah M. Buchanan¹, Shana B. Mintz¹, Weihow Hsue¹

¹Center of Animal Resources and Education, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Shana B. Mintz, DVM, DACVIM (Cardiology), Department of Clinical Sciences, smb33@cornell.edu

Co-mentor: Weihow Hsue DVM, DACVIM (Cardiology), Department of Clinical Sciences, wh446@cornell.edu

Abstract:

Background and Objectives

Although permanent pacemakers are routinely used to treat symptomatic bradyarrhythmias, long-term changes in interrogated and programmed parameters are not well characterized. The primary goal is to identify pacemaker parameters that significantly change over time. Secondary aims include comparing patient or pacemaker characteristics, such as body weight, generator brand, lead fixation type, etc., in relation to these changes and to complications.

Methods


Seventy-three client-owned dogs who received a pacemaker and had at least two subsequent interrogations were retrospectively included. Patient characteristic and pacemaker parameters were documented. Timing of recheck interrogations were binned for analysis in a linear mixed effects model. Complication frequency was compared between characteristics.

Results

Pulse width, battery life, and battery impedance significantly changed over time. Patients under seven years old had less battery life overall (P=0.0014) as well as at multiple time points. Patients with endocardial leads with active fixation had a higher amplitude threshold (P=0.0013), programed amplitude (P=0.0073), and sensitivity (P=<0.0001) compared to passive fixation, but there was no difference in the frequency of complications between types of fixation. Medtronic generators had more programming complications (P=0.0102) and required a higher programmed sensitivity (P=0.0018). A higher sensitivity (P=<0.0001) was also programmed in those with bipolar leads.

Conclusions and Clinical Importance

Some pacemaker parameters changed over time, highlighting the necessity for evaluation of the individual patient. Certain patient or pacemaker characteristics may influence pacemaker function, with lead fixation type showing particular importance but may not lead to more complications.



Alberto Oramas, DVM

ao374@cornell.edu

Institution and Location

University of Las Palmas De Gran Canaria
Cornell University, Ithaca, New York

Degree

DVM
Residency

Year

2013
2022-Present

Current Position

2nd Year Resident in Small Animal Surgery

Abstract Title:

STERNAL RECUMBENCY FOR LAPAROSCOPIC RIGHT LATERAL LIVER LOBECTOMY WITH THE USE OF TRANS-DIAPHRAGMATIC PORT. DESCRIPTION OF SURGICAL APPROACH AND CADAVERIC STUDY

Authors Names:

Alberto Oramas¹, Nicole J. Buote¹

¹Department of Clinical Sciences, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Nicole J. Buote, DVM, DACVS, MIS Founding Fellow, Department of Clinical Sciences, njb235@cornell.edu

Abstract:

Introduction:

The primary objective of this study is to evaluate the feasibility of laparoscopic right lateral liver lobectomy in canine cadavers positioned on sternal recumbency with the aid of an intrathoracic trans-diaphragmatic trocar (ITT). We hypothesized that right lateral liver lobectomies would be successfully performed with the suggested laparoscopic approach regardless of patient size.

Methods:

Ten canine cadavers euthanized for reasons unrelated to the study underwent laparoscopic right lateral liver lobectomies in sternal recumbency with an ITT. After liver lobectomy all livers were harvested. Distance from the transection to the hilus was recorded. Total surgical time, liver lobectomy surgical time, weight of the liver lobes, and complications were documented for each cadaver.

Results:

The median cadaveric weight was 19.5 kg (range 10.4-38). Median total surgical time was 34.1 minutes (range 25.1- 62.8). Median liver lobectomy surgical time was 17.8 minutes (range 7.2-42.9). The median weight of the right lateral liver lobe resected was 52.5 grams (range 15- 250). The mean distance to the hilus was 19.8 mm (STD +/- 9.4). All liver lobectomies were successfully completed.

Conclusion:

In conclusion, laparoscopic right lateral liver lobectomy is feasible in canine patients of different sizes in sternal recumbency. The use of a trans-diaphragmatic port can help to evaluate proximity to the hilus and manipulation of the liver lobes.



Shanna Wong, DVM

sw834@cornell.edu

Institution and Location

North Carolina State University, Raleigh, North Carolina
The Animal Medical Center, New York, New York
Cornell University, Ithaca, New York
Cornell University, Ithaca, New York

Degree

DVM
Internship
Sp. Internship
Residency

Year

2020
2021
2022
2022-Present

Current Position

2nd Year Resident in Anesthesia and Pain Management

Abstract Title:

AMBULATORY SACROCOCCYGEAL EPIDURAL INJECTION PROVIDES SUPERIOR POST-OPERATIVE ANALGESIA AFTER FELINE OVARIOHYSTERECTOMY (OVH)

Authors Names:

Shanna Wong¹, Richard Muto¹, Kailey Tobin¹, Stephen Parry², Manuel Martin-Flores¹, Luis Campoy¹, Robin Gleed¹, Jordyn Boesch¹

¹Department of Clinical Sciences, Cornell University, Ithaca, New York

²Cornell Statistical Consulting Unit, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Jordyn M. Boesch, DVM, PhD, DACVAA, Department of Clinical Sciences, jmb264@cornell.edu

Abstract:

An effective ambulatory sacrocoxygeal epidural technique could improve recovery of cats after OVH. We hypothesized that, in cats administered a standard anesthetic and analgesic drug protocol, an epidural injection would improve post-operative analgesia compared to the standard protocol alone. Healthy female cats undergoing OVH were assigned randomly to control or epidural groups (n = 29 each). After surgery and immediately before recovery, an electrolocation-guided sacrocoxygeal epidural injection was administered to the epidural group (0.75 mL/kg injectate volume, 0.1 mg/kg PF morphine, 1 mg/kg PF bupivacaine, remainder 0.9% saline). Using the Feline Grimace Scale (FGS), a blinded anesthesiologist assessed pain after extubation (T0) at T1, 2, 4, 6, 8, 12, and 20 hours. Whenever FGS score exceeded 3, rescue analgesia (2 mg/kg robenacoxib SC) was provided. Kaplan-Meier survival curves demonstrated a significant (log-rank test, p < 0.0001) difference between groups in time to rescue analgesia, with robenacoxib required in 24/29 cats (83%; all at T1) and 6/29 cats (21%; 4 cats at T1, 2 cats at T2) in the control and epidural groups, respectively (p < 0.0001). A linear mixed effect model showed a significant effect of the interaction of time and group; mean FGS scores (mean ± standard deviation) at T1 were 5.3 ± 2.5 and 1.8 ± 1.8 in the control and epidural groups, respectively (p < 0.0001). Adding an ambulatory sacrocoxygeal epidural injection of morphine and bupivacaine to a standard drug protocol provided superior analgesia for up to 20 hours after OVH.



Rachel P. Mandelbaum, DVM

rpm247@cornell.edu

Institution and Location

Michigan State University; East Lansing, Michigan
The Animal Medical Center; New York, New York
The Animal Medical Center; New York, New York
Cornell University, Ithaca, New York

Degree

DVM
Internship
Specialty Intern.
Residency

Year

2021
2022
2023
2023-Present

Current Position

1st Year Resident in Anesthesiology

Abstract Title:

COMPARISON OF TWO LOCATIONS FOR EXTERNAL JUGULAR VEIN CANNULATION IN A PIG MODEL

Authors Names:

Rachel P. Mandelbaum¹, Cristina de Miguel Garcia¹, Barbara Delvescovo¹, Shanna Wong¹, Julia Bledsoe¹, Kailey Tobin¹, Jordyn Boesch¹, Stephanie Hon¹

¹Department of Clinical Sciences, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Stephanie A. Hon DVM DACVAA, Department of Clinical Sciences, sar299@cornell.edu

Co-Mentor: Barbara Delvescovo, DVM, DACVIM, Department of Clinical Sciences, bd382@cornell.edu

Abstract:

Background/Rationale

Research pig models often require external jugular vein (EJV) cannulation to accomplish intricate study design elements. Although historically surgically placed, EJV cannulation via minimally invasive, ultrasound guidance techniques has been described; however, optimal placement location has not. This project aims to compare two locations for EJV cannulation based on two factors associated with successful placement: introducer insertion angle (INA - angle between introducer needle trajectory and superficial EJV wall trajectory) and maximum vessel diameter (MVD - the maximum distance measured from superficial to deep vessel wall in longitudinal plane).

Hypothesis and Scientific Design

The primary hypothesis is that the INA will be more acute in the caudal neck compared to the middle neck, the secondary hypothesis is that the MVD will be larger in the caudal neck compared to the middle neck. Eight pigs will be anesthetized and placed into standard positioning for cannulation. The caudal and middle neck will be marked via anatomical landmarks. At each location, INA and MVD will be recorded. Data will be tested for normality using a Shapiro–Wilk test and compared using a Student’s t-test or Wilcoxon rank sum test, based on normality.

Expected Outcomes

From clinical experience, the authors are consistently able to place EJV cannulas in the caudal neck, and anecdotally the difference in success between operators seems to be placement location. Cadaver dissection and initial pilot work support the hypotheses, and expected outcomes are a more acute INA and larger MVD in the caudal neck compared to the middle neck.



Corene A Bruhns, DVM

cab548@cornell.edu

Institution and Location

Cornell University, Ithaca, New York
Cornell University, Ithaca, New York

Degree

DVM
Residency

Year

2021
2022-Present

Current Position

2nd Year Resident in Medical Oncology

Abstract Title:

CHARACTERIZATION AND ASSESSMENT OF BIOLOGIC BEHAVIOR OF CANINE SPLENIC STROMAL SARCOMAS

Authors Names:

Corene A Bruhns¹, Cheryl E Balkman¹, Skylar R Sylvester¹, Andrew D Miller²

¹Department of Clinical Sciences, Cornell University, Ithaca, New York

²Department of Biomedical Sciences, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Cheryl Balkman, DVM ACVIM (SAIM, Oncology), Department of Clinical Sciences, ceb11@cornell.edu

Co-Mentor: Skylar Sylvester, DVM ACVIM (Oncology), Department of Clinical Sciences, srs289@cornell.edu


Abstract:

Background/Rationale

The diagnosis splenic stromal sarcomas (SSS) is applied to a variety of spindle cell tumors of the spleen, but there can be considerable histologic crossover with other primary splenic sarcomas. Histologic nomenclature has revised the former diagnosis of fibrohistiocytic nodules into more specific subtypes, including SSS, in the last decade. Clinical data regarding specific treatment and outcome for SSS is not well described. Developing an understanding of the tissue of origin to determine if these tumors represent distinct subtypes is clinically important. If a subset of SSS represent poorly defined counterparts of more common malignancies, such as hemangiosarcoma or histiocytic sarcoma, this significantly impacts patient prognosis, outcome, and treatment recommendations.

Hypothesis, Scientific Design, and Expected Outcomes

The hypothesis of this study is that a panel of immunohistochemical (IHC) stains applied to previously diagnosed SSS cases will alter the histologic diagnosis in a subset of cases and better characterize this group of tumors. To investigate this hypothesis, electronic medical records will be searched for cases of canine splenic masses diagnosed as SSS and fibrohistiocytic nodule(s). All cases with tissue samples identified will have IHC markers applied to confirm mesenchymal origin (vimentin) and characterize poorly-differentiated hemangiosarcoma (vWF/CD31), histiocytic sarcoma (IbA1), and leiomyosarcoma (alpha smooth muscle actin). The tumors will then be evaluated to determine if IHC alters the diagnosis, and the distribution and frequency of tumor subtypes in the population. Medical records will be reviewed to determine if any histologic characteristics impact patient outcomes, including progression free interval and overall survival.



Dahihm Kim, DVM, MS, PhD

ceb548@cornell.edu

Institution and Location

Institution and Location

Chonnam National University, Gwangju, South Korea
Seoul National University, Seoul, South Korea
Cornell University, Ithaca, New York

Degree

DVM

MS

Residency

Year

2021

2022-Present

Current Position

1st Year Resident in Laboratory Animal Medicine

Abstract Title:

CHARACTERIZATION AND ASSESSMENT OF BIOLOGIC BEHAVIOR OF CANINE SPLENIC STROMAL SARCOMAS

Authors Names:

Corene A Bruhns¹, Cheryl E Balkman¹, Skylar R Sylvester¹, Andrew D Miller²

¹Department of Clinical Sciences, Cornell University, Ithaca, New York

²Department of Biomedical Sciences, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Cheryl Balkman, DVM ACVIM (SAIM, Oncology), Department of Clinical Sciences, ceb11@cornell.edu

Co-Mentor: Skylar Sylvester, DVM ACVIM (Oncology), Department of Clinical Sciences, srs289@cornell.edu

Abstract:

Background/Rationale

The diagnosis splenic stromal sarcomas (SSS) is applied to a variety of spindle cell tumors of the spleen, but there can be considerable histologic crossover with other primary splenic sarcomas. Histologic nomenclature has revised the former diagnosis of fibrohistiocytic nodules into more specific subtypes, including SSS, in the last decade. Clinical data regarding specific treatment and outcome for SSS is not well described. Developing an understanding of the tissue of origin to determine if these tumors represent distinct subtypes is clinically important. If a subset of SSS represent poorly defined counterparts of more common malignancies, such as hemangiosarcoma or histiocytic sarcoma, this significantly impacts patient prognosis, outcome, and treatment recommendations.

Hypothesis, Scientific Design, and Expected Outcomes

The hypothesis of this study is that a panel of immunohistochemical (IHC) stains applied to previously diagnosed SSS cases will alter the histologic diagnosis in a subset of cases and better characterize this group of tumors. To investigate this hypothesis, electronic medical records will be searched for cases of canine splenic masses diagnosed as SSS and fibrohistiocytic nodule(s). All cases with tissue samples identified will have IHC markers applied to confirm mesenchymal origin (vimentin) and characterize poorly-differentiated hemangiosarcoma (vWF/CD31), histiocytic sarcoma (IbA1), and leiomyosarcoma (alpha smooth muscle actin). The tumors will then be evaluated to determine if IHC alters the diagnosis, and the distribution and frequency of tumor subtypes in the population. Medical records will be reviewed to determine if any histologic characteristics impact patient outcomes, including progression free interval and overall survival.



Victoria R. Albano, DVM

vra23@cornell.edu

Institution and Location

Cornell University, Ithaca, New York
Animal Medical Center, New York, New York
Cornell University, Ithaca, New York

Degree

DVM
Internship
Residency

Year

2021
2022
2023-Present

Current Position

2nd Year Resident in Anesthesia and Pain Management

Abstract Title:

REFINEMENT OF CONTINUOUS POSITIVE AIRWAY PRESSURE (CPAP) THERAPY IN DOGS

Authors Names:

Victoria Albano¹, Stephen Parry² Joaquin Araos ¹

¹Department of Clinical Sciences, Cornell University, Ithaca, New York

²Cornell Statistical Consulting Unit, Cornell University, Ithaca, New York

Project Mentor(s):

Mentor: Joaquin Araos, MV, PhD, DACVAA, Department of Clinical Sciences (Section of Anesthesia and Pain Management), jda246@cornell.edu

Abstract:

Postoperative respiratory dysfunction is common in dogs. Continuous positive airway pressure (CPAP) is a non-invasive ventilatory technique that increases airway diameter and functional residual capacity (FRC) leading to improved overall ventilatory function. Despite its increased use, several variables, such as fresh gas flow (FGF) rate and CPAP levels are empirically selected, even though inappropriate settings could lead to hypoventilation, hemodynamic impairment, and suboptimal FRC maximization. Therefore, this proposal aims to evaluate the impact of 1) Three rates of FGF on reinhalation; 2) Three levels of CPAP on hemodynamic stability and FRC. Eight healthy research beagles will be sedated with a propofol controlled rate infusion. For Aim 1, dogs will be administered FGF rates of 10, 15, and 20 liters per minute (Lpm), in random order, at a fixed 5cmH₂O CPAP level. The main outcomes will be the partial pressure of arterial CO₂ (PaCO₂) and the inspired pressure of CO₂ (PiCO₂) measured by arterial blood gases and side-stream capnography, respectively. For Aim 2, and after a washout period, dogs will be re-sedated, and CPAP will be randomly administered at 5, 8 and 12 cmH₂O, at a fixed oxygen flow rate of 15 Lpm. The main outcomes will be cardiac output, measured by transpulmonary thermodilution, mean arterial blood pressure (MAP), and end-expiratory lung impedance (EELI), a surrogate of FRC, measured by electrical impedance tomography. We expect to generate relevant knowledge that will inform clinicians on how to optimize and individualize the rapidly expanding therapy with CPAP.

